

ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

Gold
VICTORY[®]

170802

HERBICIDE

Contains: 50 g/litre picloram plus 100 g/litre triclopyr
both as amine salts in the form of a soluble concentrate

**Controls many broadleaf weeds in pasture,
turf and amenity areas**

PLEASE READ THE ENTIRE LABEL PRIOR TO MIXING OR USE SHAKE WELL BEFORE USE.**BEWARE:** Apply this product carefully. Very toxic to some plant species.**HAZARD CLASSIFICATION:** 6.4A, 6.9B, 9.1C, 9.2A**APPROVED HANDLER:** This product must be under the control of an approved handler during use.

See SDS for full details.

RECORD KEEPING: Records of use must be kept under certain circumstances. See SDS for full details.**WARNING:** May cause eye irritation, avoid contact with the eyes. May cause organ damage from repeated oral exposure of high doses.**FIRST AID:** If swallowed DO NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If splashed in eyes, wash out immediately with water for at least 15 minutes. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.**PERSONAL PROTECTION:** Avoid contact with skin or inhalation of spray mist. Gloves should be worn at all times. Wear eye protection when handling the concentrate. Ensure adequate ventilation when handling the concentrate. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work. Do not eat, drink or smoke while using this product.**ECOTOXIC:** Harmful to aquatic organisms. Avoid contamination of any water supply with chemical or empty container. Very toxic to the soil environment.**DISPOSAL:** Triple rinse empty container and add rinsate to the spray tank. If recycling, discard cap and deliver clean container to an Agreco depot. Alternatively crush and bury in an approved landfill. Dispose of product only by using according to the label, or at an approved landfill.**STORAGE:** Store in original container, tightly closed, away from foodstuffs. Keep out of reach of children.

Use spray solutions the same day.

SAFETY DATA SHEET: Additional information is listed in the SDS available from Lonza NZ Limited or www.lonza.com**Approved pursuant to the HSNO Act 1996,** No: HSR000555. See www.epa.govt.nz for approval conditions.**Registered pursuant to the ACVM Act 1997,** No: P007889. See www.foodsafety.govt.nz for registration conditions.**WEED SUSCEPTIBILITY** For the most effective results applications should be made during soft active growth. Spraying when plants are dormant or stressed may give variable results.**Susceptible Broadleaf Weeds Include:** Cape daisy, catsear, chickweed, clovers, cotula, cudweed, daisy, dandelion, dichondra, dock, hawkbit, hawksbeard, hydrocotyle, mallow, mayweed, nettles, onehunga weed, oxalis (creeping), parsley piert, plantain, pearlwort, penny royal, ragwort, shepherds purse, sorrel, storksbill, thistles, twincrest, wireweed.**Moderately Susceptible Weeds Include:** Buttercup, selfheal, yarrow.**Resistant Weeds Include:** All grasses and speedwells.**DIRECTIONS FOR USE****It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.****PASTURE** High Volume Gun and hose application

WEEDS	RATE mL/100 litres	COMMENTS
Ragwort		Apply as a wetting spray to full coverage while weeds are in active growth.
- Cabbage rosettes	300	
- Multicrowned plants	400	
- Flowering plants	500	
Fennel, goat's rue, hemlock, horehound, inkweed, lupin, nettles, thistles	500	
Alligator weed, Californian thistle, capeweed, field bindweed (convulvulus), docks, sorrel, and other difficult to kill broadleaf weeds.	600	

PASTURE Other application methods

EQUIPMENT	RATE	COMMENTS
Knapsack and battery powered units (e.g C-DAX) bike sprayers.	12 ml/litre water	Apply to full coverage.
Drench gun fitted with extension lance and solid cone nozzle.	1 part Victory Gold with 20 parts water (1:20). Apply 10 ml of spray mix per plant. 5 ml/plant (1:20) or use 10 mL/plant (1:40) dilution	For annual thistles and small, first year nodding thistle use only.
Mistblower	40 ml/litre water 200 ml/litre water	When using ULV attachments.
Carpet type weedwiper	1 part Victory Gold with 20 parts water (1:20)	For control of ragwort and Californian thistle. Ensure wiper remains saturated while in use. Two passes in opposite directions are recommended. Perennial specie such as Californian thistle may require repeat treatment. Drain and wash the wiper with water immediately after use.
Wick wiper	1 part Victory Gold with 1 part water (1:1)	

ANIMAL SAFETY: Poisonous plants may become more palatable after spraying, and livestock should be kept out until plants have died down.**CAUTION:** Re-Establishment of Clovers: Picloram, an active ingredient of Victory Gold is injurious to legumes. Residues in the soil may prevent early re-establishment after treatment. The persistence of residues in soil will vary with site, climate and rate of herbicide applied. It is suggested that small areas be oversown with clover 1-2 months prior to the planned sowing time and these plants inspected for growth inhibition.**TURF AND AMENITY AREAS****(Golf courses, sports fields and playing fields).**

METHOD	RATE	DIRECTIONS
Boom	4 litres per hectare	Best results are achieved by spraying actively growing young weeds. Apply at any time of the year to turf at least 12 months old.
Gun	500 ml per 100 litres	
Spot	12 ml per litre	

Speedwell Control: Add Quantum® (500 g/litre diflufenican) at 200 ml/ha (boom) or 25 ml/100 litres (gun) or 0.5 ml/litre (spot) for scrambling, turf, and field speedwell control.**NOT FOR USE IN HOME GARDENS.** For a period of 6 months following treatment, clippings from treated areas must not be used in compost heaps, used as a mulch around sensitive plants or disposed of at any garden waste recycling centre.**RAINFASTNESS:** Rain falling before the spray has dried may result in reduced efficacy.**MIXING:** Pour the correct amount of Victory Gold into the partly filled tank, and while agitating add the remainder of the water.**WARNINGS:**• **Drift:** Apply this product carefully. **Spray drift may cause serious damage to other desirable plants.** Crops susceptible to Victory Gold include but are not limited to: all brassicas, beans, flowers, fruit trees, legumes, lupins, ornamentals, potatoes, shade trees, squash, sunflowers, tomatoes, vegetables and vine crops.• **Sprayer clean-up:** To avoid injury to subsequently sprayed crops, thoroughly remove all traces of Victory Gold from mixing and spray equipment. Thoroughly clean all sprayer parts with several flushes of clean water.**RESISTANCE MANAGEMENT:** **GROUP 03 HERBICIDE** Victory Gold contains two herbicides from the pyridine carboxylic acid group with the auxin mimic mode of action. Certain weed biotypes resistant to Victory Gold and related group O3 herbicides could exist or develop due to natural genetic variation. To minimise the risk of any resistant plants dominating the weed population use strictly in accordance with label instructions. Do not use group O2 herbicides exclusively and follow resistance management strategies that may exist for particular weeds on this label. For further information see www.resistance.nzpps.org**NOTICE TO BUYER:** The manufacturer and vendors of Victory Gold warrant the formulation to contain the specified active ingredients within accepted tolerance when packed but make no warranty of any kind on the use of this material and accept no responsibility for results, whether the material has been used in accordance with directions or not.**REGISTERED TO:** Lonza NZ Limited, Hudson Road, New Plymouth, NZ.

Tel: 06-755-9234 Fax: 06-755-1174

EMERGENCY PHONE: 0800 CHEMCALL 0800 243 6225